

RESEARCH ARTICLE | JUNE 20 2013

## On an extension problem for density matrices

Eric A. Carlen; Joel L. Lebowitz; Elliott H. Lieb



*J. Math. Phys.* 54, 062103 (2013)

<https://doi.org/10.1063/1.4808218>



### Articles You May Be Interested In

On exchangeable continuous variable systems

*J. Math. Phys.* (January 2009)

Generating random density matrices

*J. Math. Phys.* (June 2011)

Lower bound on entanglement in subspaces defined by Young diagrams

*J. Math. Phys.* (January 2019)



## Special Topics Open for Submissions

[Learn More](#)

## On an extension problem for density matrices

Eric A. Carlen,<sup>1</sup> Joel L. Lebowitz,<sup>2</sup> and Elliott H. Lieb<sup>3</sup>

<sup>1</sup>*Department of Mathematics, Hill Center, Rutgers University, 110 Frelinghuysen Road, Piscataway, New Jersey 08854-8019, USA*

<sup>2</sup>*Departments of Mathematics and Physics, Hill Center, Rutgers University, 110 Frelinghuysen Road, Piscataway, New Jersey 08854-8019, USA*

<sup>3</sup>*Departments of Mathematics and Physics, Jadwin Hall, Princeton University, Washington Road, Princeton, New Jersey 08544-0001, USA*

(Received 9 February 2013; accepted 16 May 2013; published online 20 June 2013)

We investigate the problem of the existence of a density matrix  $\rho_{123}$  on a Hilbert space  $\mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \mathcal{H}_3$  with given partial traces  $\rho_{12} = \text{Tr}_3 \rho_{123}$  and  $\rho_{23} = \text{Tr}_1 \rho_{123}$ . While we do not solve this problem completely, we offer partial results in the form of some necessary and some sufficient conditions on  $\rho_{12}$  and  $\rho_{23}$ . The quantum case differs markedly from the classical (commutative) case, where the obvious necessary compatibility condition suffices, namely,  $\text{Tr}_1 \rho_{12} = \text{Tr}_3 \rho_{23}$ . © 2013 AIP Publishing LLC. [<http://dx.doi.org/10.1063/1.4808218>]

### I. INTRODUCTION

The problem considered here is closely related to the *quantum marginal problem*, on which there is an extensive literature. However, since the problem we consider involve *overlapping marginals*, results in the literature shed little light on it. We therefore introduce the problem in terms that we find natural, and postpone the discussion of the relation between our results and results on the quantum marginal problem until later in the Introduction.

Let  $\mathcal{H}_1, \mathcal{H}_2, \mathcal{H}_3$  be three finite dimensional Hilbert spaces. Let  $\rho_{12}$  be a density matrix on  $\mathcal{H}_{12} = \mathcal{H}_1 \otimes \mathcal{H}_2$  and, similarly, let  $\rho_{23}$  be a density matrix on  $\mathcal{H}_{23} = \mathcal{H}_2 \otimes \mathcal{H}_3$ . The question we ask, and which we can only partially resolve, is:

*Assuming that there is consistency of the partial traces, namely,  $\text{Tr}_1 \rho_{12} = \rho_2 = \text{Tr}_3 \rho_{23}$ , what are necessary and sufficient conditions for the existence of a density matrix  $\rho_{123}$  on  $\mathcal{H}_{123} = \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \mathcal{H}_3$  such that  $\text{Tr}_1 \rho_{123} = \rho_{12}$  and  $\text{Tr}_3 \rho_{123} = \rho_{23}$ ?*

This question goes back at least to Werner,<sup>17</sup> who proved the existence of consistent pairs for which no extension exists. A simple example is given later in this Introduction.

This is an obvious question to ask in several contexts, e.g., Ref. 9. We note the fact that in classical statistical mechanics the answer is that an extension always exists, and there is a simple formula for it: Given probability densities  $\rho_{12}(x, y)$  and  $\rho_{23}(y, z)$  with

$$\int \rho_{12}(x, y) d\mu_1(x) = \int \rho_{23}(y, z) d\mu_3(z) =: \rho_2(y),$$

we may define

$$\rho_{123}(x, y, z) = \frac{\rho_{12}(x, y)\rho_{23}(y, z)}{\rho_2(y)}. \quad (1.1)$$

If the sample spaces over which  $x, y$ , and  $z$  range are finite sets, these densities may be identified with diagonal density matrices in a natural way, embedding the discrete classical probability space problem into the quantum mechanical problem.

This extension is not unique, in general, but among all extensions, this one has the maximum entropy, namely,  $S_{12} + S_{23} - S_2$ . The maximality is a consequence of the Strong Subadditivity of

Entropy (SSA) which says that for any extension

$$S_{12} + S_{23} \geq S_{123} + S_2.$$

The classical SSA inequality is relatively easy to prove (as opposed to its quantum version); see Ref. 11. The principle behind the construction is *conditioning*: Note that for random variables  $Y$  and  $Z$  with joint density  $\rho_{23}(y, z)$ ,  $\rho_{23}(Y, z)/\rho_2(Y)$  is the conditional density for  $Z$  given  $Y$ .

This construction of extensions by conditioning can be generalized to arbitrarily many factors. Given  $\rho_{j,j+1}$ ,  $j = 1, \dots, N$ , such that

$$\int \rho_{j-1,j}(x_{j-1}, x_j) d\mu_{j-1}(x_{j-1}) = \int \rho_{j,j+1}(x_j, x_{j+1}) d\mu_{j+1}(x_{j+1}) =: \rho_j(x_j)$$

define

$$\rho_{1,\dots,N}(x_1, \dots, x_N) = \rho_{1,2}(x_1, x_2) \prod_{j=2}^{N-1} \frac{\rho_{j,j+1}(x_j, x_{j+1})}{\rho_j(x_j)}.$$

Again this extension is the maximum entropy extension by an iterated application of SSA.

However, the construction (1.1), being based on conditional probabilities, does not generalize to the quantum case, where the notion of “conditioning” has no obvious meaningful analog. In general, even for consistent density matrices,

$$\rho_{12}\rho_2^{-1}\rho_{23} \quad (1.2)$$

need not be Hermitian, much less positive definite. While

$$R = \exp[\log \rho_{12} + \log \rho_{23} - \log \rho_2]$$

is Hermitian and positive definite, its partial traces will not equal the desired reduced density matrices, even after renormalization, which would be required since the trace,  $\text{Tr}_{123}R$ , is never greater than 1, and, generally, is less than 1. This fact follows from the triple Golden-Thompson inequality<sup>10</sup>

$$\text{Tr}_{123} R \leq \text{Tr}_2 \text{Tr}_{13} \int_0^\infty \rho_{12} \frac{1}{t + \rho_2} \rho_{23} \frac{1}{t + \rho_2} dt = \text{Tr}_2 \int_0^\infty \rho_2 \frac{1}{t + \rho_2} \rho_2 \frac{1}{t + \rho_2} dt = \text{Tr}_2 \rho_2 = 1.$$

One situation in which the classical formula (1.2) is valid occurs when  $\rho_{12}$  and  $\rho_{23}$  commute and are consistent. This implies that  $\rho_2$  commutes with  $\rho_{12}$  and  $\rho_{23}$ . (To see, note that  $\rho_2\rho_{23} = \text{Tr}_1(\rho_{12}\rho_{23}) = \text{tr}_1(\rho_{23}\rho_{12}) = \rho_{23}\rho_2$ , and likewise for  $\rho_2\rho_{12} = \rho_{12}\rho_2$ .)

We note that if  $\rho_{12}$  is pure and  $\rho_{12} = \text{Tr}_3 \rho_{123}$ , then  $\rho_{123} = \rho_{12} \otimes \rho_3$ . Hence  $\rho_{23}$  must be a product when there is an extension  $\rho_{123}$ . This provides simple examples of compatible pairs with no extension. For example, take  $\rho_{12}$  to be pure and  $\rho_{23}$  not to be a product, but consistent with  $\rho_{12}$ .

In Sec. II, Theorem 2.1 generalizes this observation by identifying a class of non-pure states  $\rho_{12}$ , which can only be extended by product states  $\rho_{12} \otimes \rho_3$ .

Motivated by these examples, and the obvious difference between the classical and quantum cases, we believe that a proper understanding of this problem will lead to a clearer understanding of entanglement and quantum information theory.

## A. The quantum marginal problem

The quantum marginal problem, in its simplest form, is the following: Given density matrices  $\rho_1$  and  $\rho_2$ , on  $\mathcal{H}_1$  and  $\mathcal{H}_2$ , respectively, let  $\mathcal{C}(\rho_1, \rho_2)$  denote the set of all density matrices  $\rho_{12}$  on  $\mathcal{H}_1 \otimes \mathcal{H}_2$  such that

$$\text{Tr}_1 \rho_{12} = \rho_2 \quad \text{and} \quad \text{Tr}_2 \rho_{12} = \rho_1. \quad (1.3)$$

Here,  $\mathcal{C}(\rho_1, \rho_2)$  is the set of *quantum couplings* of  $\rho_1$  and  $\rho_2$ .

Note that  $\mathcal{C}(\rho_1, \rho_2)$  is never empty; it always contains  $\rho_1 \otimes \rho_2$ . It is also evidently convex and compact, and hence it is the convex hull of its extreme points. Characterizations of the extreme points have been given by Parthasarathy<sup>14</sup> and Rudolph.<sup>16</sup>

Other research has focused on the relations between the spectra of  $\rho_1$ ,  $\rho_2$ , and  $\rho_{12}$  that characterize the set of triples of density matrices satisfying (1.3). For results in this direction see the recent papers of Klyachko,<sup>8</sup> and Christandl *et al.*,<sup>4</sup> and references therein.

The general quantum marginal problems concern states on the product of arbitrarily many Hilbert spaces, and the marginals obtained by taking any combination of partial traces – for instance, the partial traces  $\rho_{12}$  and  $\rho_{23}$  of  $\rho_{123}$  as in our problem. Most results pertain to the case of non-overlapping marginals, in contrast to the problem considered here. An exception is the paper by Osborne.<sup>13</sup> He uses the SSA of the von Neumann entropy<sup>12</sup> to prove an upper bound on the number of *orthogonal pure states*  $\rho_{123}$  such that  $\text{Tr}_1 \rho_{123} = \rho_{23}$  and  $\text{Tr}_3 \rho_{123} = \rho_{12}$ . We shall also employ entropy bounds, but are mainly concerned with the existence, or not, of *mixed-state* extensions  $\rho_{123}$  of compatible pairs  $\rho_{12}$  and  $\rho_{23}$ . For other work on overlapping marginals see the papers of Butterly *et al.*<sup>2</sup> and Hall,<sup>6</sup> who investigate necessary and sufficient conditions for the existence of an extension when one is given *all* of the proper marginals (e.g., for three spaces one has to be given  $\rho_{12}$ ,  $\rho_{13}$ , and  $\rho_{23}$ .)

It is natural to use entropy in this investigation, and some early results on the two-space problem are given in terms of entropy. For example, if  $\rho_{12}$  is a pure state, then its partial traces  $\rho_1$  and  $\rho_2$  have the same non-zero spectrum.<sup>1</sup> Conversely, if  $\rho_1$  and  $\rho_2$  are two density matrices having the same non-zero spectrum, then there is a pure state  $\rho_{12}$  satisfying (1.3). To construct it, assume without loss of generality that  $\mathcal{H}_1$  and  $\mathcal{H}_2$  have the same dimension. There is a unitary  $U$  such that  $U\rho_2U^* = \rho_1$ . Define  $\Psi := U\sqrt{\rho_2}$ , where the matrix on the right is regarded as a vector in  $\mathcal{H}_1 \otimes \mathcal{H}_2$ . Then  $\rho_{12} = |\Psi\rangle\langle\Psi|$  is such a pure state.

If  $\rho_1$  and  $\rho_2$  do not have the same spectrum, then the set  $\mathcal{C}(\rho_1, \rho_2)$  cannot contain any pure state. It is natural then to ask for the least entropy element of  $\mathcal{C}(\rho_1, \rho_2)$ . Since the entropy is concave, the minimum will be attained at an extreme point, and the results of Refs. 14 and 16 are relevant, and lead easily to the minimizer in specific cases. However, a well-known inequality already provides a sharp *a priori* lower bound for this minimum entropy coupling.

The *Araki-Lieb Triangle inequality*<sup>1</sup> says that when (1.3) is satisfied

$$S_{12} \geq |S_1 - S_2|. \quad (1.4)$$

For an interesting discussion of entropy inequalities for a general *classical* marginal problem, see Ref. 5.

## II. NECESSARY CONDITIONS FOR THE EXISTENCE OF AN EXTENSION

Strong Subadditivity of Entropy<sup>12</sup> (see also Ref. 11) provides us with two necessary conditions for an extension to exist: Let  $\rho_{123}$  be an extension of  $\rho_{12}$  and  $\rho_{23}$ . Then, discarding the positive term  $S_{123}$  in SSA,

$$S_{12} + S_{23} \geq S_2. \quad (2.1)$$

Classically, one has monotonicity of the entropy, meaning  $S_{12} \geq S_2$  and  $S_{23} \geq S_2$  so that (2.1) hold classically with  $2S_2$  on the right side. However, quantum mechanically, equality can hold in (2.1): Let  $\rho_{12}$  be a purification of  $\rho_2$ , i.e., a pure state  $|\Psi\rangle\langle\Psi|$  on  $\mathcal{H}_1 \otimes \mathcal{H}_2$  such that  $\text{Tr}_2(|\Psi\rangle\langle\Psi|) = \rho_1$ . Such purifications always exist. It is well known, and easy to see from the definition, that the set of all possible purifications  $\Psi$  of  $\rho_1$  is the set of all operators  $\sqrt{\rho_1}U$ , regarded as vectors in  $\mathcal{H}_1 \otimes \mathcal{H}_2$ , where  $U$  is a partial isometry from a subspace of  $\mathcal{H}_2$  onto the range of  $\rho_1$ . Let  $\rho_{23}$  be the tensor product of  $\rho_2$  and a pure state. Then  $S_{12} = 0$  and  $S_{23} = S_2$ .

A second form of SSA<sup>12</sup> leads to a sharper necessary condition: Again assume that  $\rho_{123}$  is an extension of  $\rho_{12}$  and  $\rho_{23}$ . Then

$$S_{12} + S_{23} \geq S_1 + S_3. \quad (2.2)$$

As we now explain, whenever (2.2) is satisfied by any consistent pair of density matrices  $\rho_{12}$  and  $\rho_{23}$  (not necessarily possessing a common extension), then (2.1) is automatically satisfied. To see

this, note that by (1.4),  $S_2 - S_1 \leq S_{12}$  and  $S_2 - S_3 \leq S_{23}$ . Adding these inequalities, we obtain

$$2S_2 \leq S_{12} + S_{23} + S_1 + S_3.$$

By (2.2), the right side is no greater than  $2(S_{12} + S_{23})$ , which yields (2.1).

The density matrices for which there is equality in the triangle inequality have a particular structure: Let  $m$  and  $n$  be positive integers, and let  $\{\lambda_1, \dots, \lambda_m\}$  and  $\{\mu_1, \dots, \mu_n\}$  be sets of positive numbers with  $\sum_{j=1}^m \lambda_j = \sum_{k=1}^n \mu_k = 1$ . Then, as shown in Ref. 3, there exists a density matrix  $\rho_{12}$  such that the non-zero eigenvalues of  $\rho_{12}$  are  $\{\lambda_1, \dots, \lambda_m\}$ , the non-zero eigenvalues of  $\rho_2$  are  $\{\mu_1, \dots, \mu_n\}$ , and the non-zero eigenvalues of  $\rho_1$  are the numbers  $\{\lambda_j \mu_k : 1 \leq j \leq m, 1 \leq k \leq n\}$ . For any such  $\rho_{12}$ , it is evident that  $S_{12} = S_1 - S_2$ . Moreover, as shown in Ref. 3, whenever  $S_{12} = S_1 - S_2$ , the spectra of  $\rho_{12}$ ,  $\rho_1$ , and  $\rho_2$  are related in this way.

The following theorem generalizes the observation that pure states  $\rho_{12}$  may only be extended by product states  $\rho_{12} \otimes \rho_3$ . When  $\rho_{12}$  is pure,  $0 = S_{12} = S_2 - S_1$ .

**Theorem 2.1.** *Let  $\rho_{12}$  be a density matrix such that*

$$S_{12} = S_1 - S_2. \quad (2.3)$$

*Then  $\rho_{12}$  and  $\rho_{23}$  have a common extension if and only if  $\rho_{23} = \rho_2 \otimes \rho_3$ .*

*Proof.* Using (2.3) in (2.2) we obtain

$$S_1 - S_2 + S_{23} \geq S_3 + S_1,$$

which reduces to  $S_{23} \geq S_2 + S_3$ . By the subadditivity of the entropy, this means that  $S_{23} = S_2 + S_3$ , and so  $\rho_{23} = \rho_2 \otimes \rho_3$ .  $\square$

In Sec. IV, we give an example in which (2.2) is satisfied, but there is no common extension.

### III. SUFFICIENT CONDITIONS FOR THE EXISTENCE OF AN EXTENSION

We do not have any very general sufficient conditions for compatible pairs to possess a common extension. One general positive statement that can be made is the following.

**Theorem 3.1.** *Let  $\rho_{12}$  and  $\rho_{23}$  be a compatible pair of density matrices that possess a common extension  $\rho_{123}$  that is positive definite. Let  $\|\cdot\|$  denote the trace norm. Then there is an  $\epsilon > 0$ , depending on the dimensions and the smallest eigenvalue of  $\rho_{123}$ , such that if  $\tilde{\rho}_{12}$  and  $\tilde{\rho}_{23}$  is another compatible pair on the same spaces and*

$$\|\rho_{12} - \tilde{\rho}_{12}\| + \|\rho_{23} - \tilde{\rho}_{23}\| < \epsilon, \quad (3.1)$$

*then  $\tilde{\rho}_{12}$  and  $\tilde{\rho}_{23}$  possess common extension.*

*Proof.* Define

$$\tilde{\rho}_{123} = \rho_{123} + [\tilde{\rho}_{12} - \rho_{12}] \otimes \tilde{\rho}_3 + \tilde{\rho}_1 \otimes [\tilde{\rho}_{23} - \rho_{23}] + \tilde{\rho}_1 \otimes [\rho_2 - \tilde{\rho}_2] \otimes \tilde{\rho}_3.$$

It is easily checked that  $\text{Tr}_1 \tilde{\rho}_{123} = \tilde{\rho}_{23}$  and  $\text{Tr}_3 \tilde{\rho}_{123} = \tilde{\rho}_{12}$ . Furthermore,  $\tilde{\rho}_{123}$  is self-adjoint, and under the condition (3.1) is positive when  $\epsilon$  is chosen sufficiently small.  $\square$

**Remark 3.2.** Of course, in this finite dimensional setting, the trace norm could be replaced by any other norm. Also, suppose that  $\rho_{12}$  is positive definite, and  $\rho_{23} = \rho_2 \otimes \rho_3$  with  $\rho_3$  positive definite. In this case,  $\rho_{12}$  and  $\rho_{23}$  are compatible and have the positive-definite common extension  $\rho_{12} \otimes \rho_3$ . The theorem says that any compatible pair that is a small perturbation of  $\rho_{12}$  and  $\rho_{23}$  has a common extension. However, as Theorem 2.1 shows, the requirement of positive definiteness cannot be dropped.

Let  $\rho_{12}$  and  $\rho_{23}$  be a consistent pair of density matrices, both of which are separable. Recall that a bipartite density matrix  $\rho_{12}$  is *finitely separable* if it is a convex combination of product states; i.e.,

if

$$\rho_{12} = \sum_{j=1}^N \lambda_j \sigma_1^{(j)} \otimes \tau_2^{(j)},$$

where each  $\sigma_1^{(j)}$  is a density matrix on  $\mathcal{H}_1$ , each  $\tau_2^{(j)}$  is a density matrix on  $\mathcal{H}_2$  and, crucially, each  $\lambda_j > 0$ . The set of separable density matrices is the closure of the set of finitely separable density matrices.

Separable density matrices are often viewed as being a classical ensemble of product states. As such, separable states are free of *quantum correlations* among observables on  $\mathcal{H}_1$  and  $\mathcal{H}_2$ . Separable bipartite density matrices do, indeed, behave much more like classical joint probability distributions. For example, if  $\rho_{12}$  is separable, then (see, e.g., Ref. 3),

$$S_{12} \geq \max\{S_1, S_2\},$$

and thus when both  $\rho_{12}$  and  $\rho_{23}$  are separable and compatible,  $S_{12} + S_{23} \geq S_1 + S_3$ , and thus the condition (2.2), which is necessary for a common extension to exist, is always satisfied.

One might therefore, hope that a common extension  $\rho_{123}$  would exist whenever  $\rho_{12}$  and  $\rho_{23}$  are compatible and both are separable. In Sec. IV, we show that this is not the case: Further conditions must be imposed to ensure the existence of an extension. Here is one case in which a common extension does exist.

Suppose that

$$\rho_{12} = \sum_{j=1}^n \lambda_j \rho^{(j)} \otimes \sigma^{(j)} \quad \text{and} \quad \rho_{23} = \sum_{j=1}^n \lambda_j \sigma^{(j)} \otimes \tau^{(j)}, \quad (3.2)$$

where the  $\rho^{(j)}$ ,  $\sigma^{(j)}$ , and  $\tau^{(j)}$  are density matrices and the  $\lambda_j$  are positive numbers with  $\sum_{j=1}^n \lambda_j = 1$ . Then

$$\rho_{123} := \sum_{j=1}^n \lambda_j \rho^{(j)} \otimes \sigma^{(j)} \otimes \tau^{(j)} \quad (3.3)$$

is a common extension.

While this example is based on a strong assumption, namely, that the weights and the factors on  $\mathcal{H}_2$  coincide, we shall show in Sec. IV that consistency and separability is not enough.

However, as we now explain, there is a more general version of this construction using coherent states. Good references on coherent states are Refs. 7 and 15.

Let  $\mathcal{H}$  be any  $d$  dimensional Hilbert space. Define  $J = (d - 1)/2$ . Then there is an irreducible representation of  $SU(2)$  on  $\mathcal{H}$ . Associated with this representation is a family of pure states  $|\Omega\rangle\langle\Omega|$  on  $\mathcal{H}$ , parameterized by a point  $\Omega$  in the unit sphere,  $S^2$ . Given any self-adjoint operator  $A$  on  $\mathcal{H}$ , there is a function  $\hat{a}(\Omega)$  on  $S^2$  such that

$$A = \int_{S^2} d\Omega \hat{a}(\Omega) |\Omega\rangle\langle\Omega|, \quad (3.4)$$

where the integration is with respect to the uniform probability measure on  $S^2$ . The function  $\hat{a}(\Omega)$  is *unique* if we further require that  $\hat{a}(\Omega)$  be a spherical harmonic of degree no higher than  $2J$ . (See p. 33 and Eq. (4.10) of Ref. 7.) This function  $\hat{a}(\Omega)$  is called the *upper symbol* of the operator  $A$ . The upper symbol of  $A$  need not be a non-negative function on  $S^2$  even if  $A$  is positive definite. However, if  $\hat{a}$  is non-negative, then the operator  $A$  defined by (3.4) is positive semidefinite.

Now consider a bipartite density matrix  $\rho_{12}$  on  $\mathcal{H}_1 \otimes \mathcal{H}_2$  where the dimension of  $\mathcal{H}_j = d_j$  for  $j = 1, 2$ . Suppose that it has a representation

$$\rho_{12} = \int_{S^2} d\Omega_2 \int_{S^2} d\Omega_1 \tilde{\rho}_{12}(\Omega_1, \Omega_2) (|\Omega_1\rangle\langle\Omega_1| \otimes |\Omega_2\rangle\langle\Omega_2|)$$

with  $\tilde{\rho}_{12}(\Omega_1, \Omega_2)$ , a non-negative function on  $S^2 \times S^2$  that is a spherical harmonic of degree  $d_j - 1$  in  $\Omega_j, j = 1, 2$ .

Then taking the trace over  $\mathcal{H}_1$ , we find

$$\rho_2 := \text{Tr}_1 \rho_{12} = \int_{S^2} d\Omega_2 \left[ \int_{S^2} d\Omega_1 \tilde{\rho}_{12}(\Omega_1, \Omega_2) \right] |\Omega_2\rangle\langle\Omega_2|,$$

and evidently  $\int_{S^2} d\Omega_1 \tilde{\rho}_{12}(\Omega_1, \Omega_2)$  has the degree  $d_2 - 1$  in  $\Omega_2$ , which ensures that it is the unique upper symbol of  $\rho_2$  of minimal degree.

Likewise, if  $\rho_{23}$  is a density matrix on  $\mathcal{H}_2 \otimes \mathcal{H}_3$  with the dimension of  $\mathcal{H}_3 = d_j$ , and if  $\rho_{23}$  is compatible with  $\rho_{12}$  and, moreover,

$$\rho_{23} = \int_{S^2} d\Omega_2 \int_{S^2} d\Omega_3 \tilde{\rho}_{23}(\Omega_2, \Omega_3) (|\Omega_2\rangle\langle\Omega_2| \otimes |\Omega_3\rangle\langle\Omega_3|)$$

with  $\tilde{\rho}_{23}(\Omega_2, \Omega_3)$  being a non-negative function on  $S^2 \times S^2$  that is a spherical harmonic of degree  $d_j - 1$  in  $\Omega_j, j = 2, 3$ ,  $\int_{S^2} d\Omega_3 \tilde{\rho}_{23}(\Omega_2, \Omega_3)$  is the upper symbol of  $\rho_2$ , and thus quantum compatibility of  $\rho_{12}$  and  $\rho_{23}$  implies the classical compatibility of the probability densities  $\tilde{\rho}_{12}(\Omega_1, \Omega_2)$  and  $\tilde{\rho}_{23}(\Omega_2, \Omega_3)$ . Hence the classical prescription may be used to define

$$\tilde{\rho}_{123}(\Omega_1, \Omega_2, \Omega_3) := \frac{\rho_{12}(\Omega_1, \Omega_2) \rho_{23}(\Omega_2, \Omega_3)}{\rho_2(\Omega_2)},$$

and it is then the case that

$$\rho_{123} := \int_{S^2} d\Omega_1 \int_{S^2} d\Omega_2 \int_{S^2} d\Omega_3 \tilde{\rho}_{123}(\Omega_1, \Omega_2, \Omega_3) (|\Omega_1\rangle\langle\Omega_1| \otimes |\Omega_2\rangle\langle\Omega_2| \otimes |\Omega_3\rangle\langle\Omega_3|)$$

is positive semidefinite, and is a common extension of  $\rho_{12}$  and  $\rho_{23}$ .

#### IV. SEPARABLE COMPATIBLE PAIRS WITH NO EXTENSION

In this section, we give examples of compatible pairs  $\rho_{12}$  and  $\rho_{23}$  that do not have an extension, but nevertheless satisfy the necessary condition (2.2). Moreover, in these examples both  $\rho_{12}$  and  $\rho_{23}$  will be separable. As we have remarked above, one might expect the extension problem to simplify in the presence of separability. This is not the case, as the following examples show. For simplicity, our examples will be on  $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ , but of course may be embedded into higher dimensional spaces.

*Lemma 4.1.* Let  $\rho_2$  be a non-pure density matrix on  $\mathbb{C}^2$ ; i.e., the rank of  $\rho_2$  is 2. Let  $\{\psi_1, \psi_2\}$  be an orthonormal basis of  $\mathbb{C}^2$  consisting of eigenvectors of  $\rho_2$ . Let  $\rho_2 \psi_j = \mu_j \psi_j, j = 1, 2$ . Then there exist two unit vectors  $\phi_1$  and  $\phi_2$  in  $\mathbb{C}^2$  and positive numbers  $v_1$  and  $v_2$  such that

$$\sum_{j=1}^2 \mu_j |\psi_j\rangle\langle\psi_j| = \rho_2 = \sum_{j=1}^2 v_j |\phi_j\rangle\langle\phi_j|, \quad (4.1)$$

and such that the four vectors  $\psi_1, \psi_2, \phi_1$ , and  $\phi_2$  are pairwise linearly independent.

*Proof.* If  $\rho_2$  is  $\frac{1}{2}\mathbb{1}$ , we may take  $\{\psi_1, \psi_2\}$  to be any orthonormal basis of  $\mathbb{C}^2$ , and then choose  $\{\phi_1, \phi_2\}$  to be any other orthonormal basis such that  $\phi_1$  is not proportional to either  $\psi_1$  or  $\psi_2$ . Then the four vectors  $\phi_1, \phi_2, \psi_1, \psi_2$  are pairwise linearly independent. In this case, we take  $\mu_j = v_j = \frac{1}{2}$  for  $j = 1, 2$ .

Next, assume  $\rho_2$  has distinct eigenvalues  $\mu_1 > \mu_2$ . Let  $\{\psi_1, \psi_2\}$  be an orthonormal basis consisting of eigenvectors of  $\rho_2$ . Let  $\phi_1$  be any unit vector that is not proportional to either  $\psi_1$  or  $\psi_2$ . Then there is a unique largest number  $v_1 > 0$ , so that  $\rho_2 - v_1 |\phi_1\rangle\langle\phi_1|$  is positive semidefinite. Thus,  $\rho_2 - v_1 |\phi_1\rangle\langle\phi_1|$  is rank one, and hence

$$\rho_2 - v_1 |\phi_1\rangle\langle\phi_1| = v_2 |\phi_2\rangle\langle\phi_2|$$

for some unit vector  $\phi_2$  that is not proportional to  $\phi_1$  since  $\rho_2$  has rank 2. Likewise,  $\phi_2$  is not an eigenvector of  $\rho_2$ , since by the non-degeneracy, this would force  $\phi_1$  to be an eigenvector, which it is not.  $\square$

**Theorem 4.2.** *Let  $\rho_2$  be a non-pure density matrix on  $\mathbb{C}^2$ . Then there exist separable density matrices  $\rho_{12}$  and  $\rho_{23}$ , extending  $\rho_2$ , and such that (2.2) is satisfied (with strict inequality), but such that  $\rho_{12}$  and  $\rho_{23}$  have no common extension  $\rho_{123}$ .*

*Proof.* Let  $\{\psi_1, \psi_2\}$  and  $\{\phi_1, \phi_2\}$  be sets of unit vectors such that (1.3) is satisfied, and where  $\{\psi_1, \psi_2\}$  is an orthonormal basis of  $\mathbb{C}^2$  consisting of eigenvectors of  $\rho$ .

Let  $\{\chi_1, \chi_2\}$  be any orthonormal basis of  $\mathbb{C}^2$ , and let  $\{\eta_1, \eta_2\}$  be any linearly independent set of unit vectors in  $\mathbb{C}^2$ . Define

$$\rho_{12} = \sum_{j=1}^2 \mu_j |\eta_j \otimes \psi_j\rangle \langle \eta_j \otimes \psi_j| \quad \text{and} \quad \rho_{23} = \sum_{j=1}^2 v_j |\phi_j \otimes \chi_j\rangle \langle \phi_j \otimes \chi_j|. \quad (4.2)$$

Then evidently  $\text{Tr}_1 \rho_{12} = \text{Tr}_3 \rho_{23} = \rho_2$ . Next, for any vector  $(z, w) \in \mathbb{C}^2$ , let  $(z, w)^\perp = (-\bar{w}, \bar{z})$  so that  $(z, w)^\perp$  is orthogonal to  $(w, z)$ .

Suppose a common extension  $\rho_{123}$  does exist. Then  $\eta_1^\perp \otimes \psi_1$  is in the nullspace of  $\rho_{12}$ , and hence

$$\sum_{j=1}^2 \langle \eta_1^\perp \otimes \psi_1 \otimes \chi_j, \rho_{123} \eta_1^\perp \otimes \psi_1 \otimes \chi_j \rangle = \langle \eta_1^\perp \otimes \psi_1, \rho_{12} \eta_1^\perp \otimes \psi_1 \rangle = 0.$$

Therefore, since  $\rho_{123}$  is positive, both  $\eta_1^\perp \otimes \psi_1 \otimes \chi_1$  and  $\eta_1^\perp \otimes \psi_1 \otimes \chi_2$  are in the nullspace of  $\rho_{123}$ .

Since  $\eta_2^\perp \otimes \psi_2$  is in the null space of  $\rho_{12}$ , the same argument shows that both  $\eta_2^\perp \otimes \psi_2 \otimes \chi_1$  and  $\eta_2^\perp \otimes \psi_2 \otimes \chi_2$  are in the nullspace of  $\rho_{123}$ . Thus, the four vectors

$$\eta_1^\perp \otimes \psi_1 \otimes \chi_j \quad \text{and} \quad \eta_2^\perp \otimes \psi_2 \otimes \chi_j, \quad j = 1, 2$$

belong to the nullspace of  $\rho_{123}$ .

Likewise, both  $\phi_1^\perp \otimes \chi_1$  and  $\phi_2^\perp \otimes \chi_2$  belong to the nullspace of  $\rho_{23}$ . Arguing as above, we see that the four vectors

$$\eta_j^\perp \otimes \phi_i^\perp \otimes \chi_1 \quad \text{and} \quad \eta_j^\perp \otimes \phi_i^\perp \otimes \chi_2, \quad j = 1, 2$$

belong to the nullspace of  $\rho_{123}$ .

Define the unit vectors

$$\Psi_1 = \eta_1^\perp \otimes \psi_1 \otimes \chi_1, \quad \Psi_2 = \eta_1^\perp \otimes \psi_1 \otimes \chi_2, \quad \Psi_3 = \eta_2^\perp \otimes \psi_2 \otimes \chi_1, \quad \text{and} \quad \Psi_4 = \eta_2^\perp \otimes \psi_2 \otimes \chi_2, \quad (4.3)$$

and the vectors

$$\Phi_1 = \eta_1^\perp \otimes \phi_1^\perp \otimes \chi_1, \quad \Phi_2 = \eta_1^\perp \otimes \phi_2^\perp \otimes \chi_2, \quad \Phi_3 = \eta_2^\perp \otimes \phi_1^\perp \otimes \chi_1, \quad \text{and} \quad \Phi_4 = \eta_2^\perp \otimes \phi_2^\perp \otimes \chi_2. \quad (4.4)$$

These 8 vectors are in the nullspace of  $\rho_{123}$  under the hypotheses imposed so far.

Now let us temporarily impose the additional hypothesis that  $\{\eta_1, \eta_2\}$  is orthonormal. Consequently,  $\{\eta_1^\perp, \eta_2^\perp\}$  is orthonormal.

We claim that the 8 unit vectors  $\Psi_j, \Phi_j, j = 1, 2, 3, 4$ , are linearly independent. To see this, note that under our additional hypothesis,  $\{\Psi_1, \Psi_2, \Psi_3, \Psi_4\}$  and  $\{\Phi_1, \Phi_2, \Phi_3, \Phi_4\}$  are orthonormal, and  $\langle \Psi_i, \Phi_j \rangle = \delta_{i,j}$ . Therefore, if

$$\sum_{j=1}^4 a_j \Psi_j + \sum_{j=1}^4 b_j \Phi_j = 0,$$

$$(a_j \Psi_j + b_j \Phi_j) = 0 \quad \text{for each} \quad j = 1, 2, 3, 4.$$



However, since the vectors  $\psi_1$  and  $\phi_1$  are linearly independent, so are the vectors  $\psi_1^\perp$  and  $\phi_2 = \phi_1^\perp$ . Thus  $\Psi_1$  and  $\Phi_1$  are linearly independent, and hence  $a_1 = b_1 = 0$ .

The same sort of argument shows that  $a_j = b_j = 0$  for each  $j = 1, 2, 3, 4$ . Thus the eight vectors  $\Psi_j, j = 1, 2, 3, 4$  and  $\Phi_j, j = 1, 2, 3, 4$  are linearly independent and in the nullspace of  $\rho_{123}$ . Since the dimension of  $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$  is 8, this means  $\rho_{123} = 0$ , which is impossible.

Therefore, the compatible pair of density matrices  $\rho_{12}$  and  $\rho_{23}$  defined by (4.2) has no common extension.

Next, observe from (4.2) that

$$\rho_1 = \sum_{j=1}^2 \mu_j |\eta_j\rangle\langle\eta_j|.$$

Since  $\eta_1$  and  $\eta_2$  are orthogonal, and since  $\eta_1 \otimes \psi_1$  and  $\eta_2 \otimes \psi_2$  are orthogonal

$$S_1 = S_{12} = - \sum_{j=1}^2 \mu_j \log \mu_j.$$

In the same manner we see that

$$S_3 = S_{23} = - \sum_{j=1}^2 v_j \log v_j.$$

Thus we have

$$S_{12} + S_{23} = S_1 + S_3.$$

Finally, making a sufficiently small change in  $\{\eta_1, \eta_2\}$ , we may arrange that this set of unit vectors is no longer orthogonal, but that the 8 unit vectors  $\Psi_j, \Phi_j, j = 1, 2, 3, 4$  defined in (4.3) and (4.4) are still linearly independent. Then the compatible pair  $\rho_{12}$  and  $\rho_{23}$  defined by (4.2) has no common extension. It is still true that  $S_{23} = S_3$ . Also, since  $\{\psi_1, \psi_2\}$  is orthonormal,  $\{\eta_1 \otimes \psi_1, \eta_2 \otimes \psi_2\}$  is orthonormal, and so  $S_{12} = - \sum_{j=1}^2 \mu_j \log \mu_j$ . However,  $\rho_1 = \sum_{j=1}^2 \mu_j |\eta_j\rangle\langle\eta_j|$  and since  $\{\eta_1, \eta_2\}$  is not orthogonal,

$$S_1 < - \sum_{j=1}^2 \mu_j \log \mu_j = S_{12}.$$

Therefore,  $S_{12} + S_{23} > S_1 + S_2$ . □

*Remark 4.3.* While the condition that  $\rho_1$  and  $\rho_2$  have the same non-zero spectrum is necessary and sufficient to ensure that there exists a common purification of  $\rho_1$  and  $\rho_2$ , the above construction shows that the condition that the nonzero spectrum of  $\rho_{12}$  equals that of  $\rho_3$ , and that the nonzero spectrum of  $\rho_{23}$  equals that of  $\rho_1$ , together with compatibility, does not ensure that there exists a common purification of  $\rho_{12}$  and  $\rho_{23}$ . Indeed, let  $\{\eta_1, \eta_2\}$ ,  $\{\psi_1, \psi_2\}$ ,  $\{\phi_1, \phi_2\}$ , and  $\{\chi_1, \chi_2\}$  be four orthonormal bases of  $\mathbb{C}^2$  having no vectors (or their opposites) in common. Then with  $\mu_j = v_j = 1/2, j = 1, 2$ , (4.2) defines a compatible pair of density matrices with  $\rho_2 = \frac{1}{2}I$ . The non-zero spectrum of  $\rho_{12}, \rho_{23}, \rho_1$ , and  $\rho_3$  is  $\{1/2, 1/2\}$ , and yet there is no common extension of  $\rho_{12}$  and  $\rho_{23}$ , as the proof of Theorem 4.2 shows.

## ACKNOWLEDGMENTS

This work is partially supported by U.S. National Science Foundation Grant Nos. DMS 1201354 (E.A.C.), DMR 113501 and AFOSR FA9550-10-1-0131 (J.L.L.), and PHY 0965859 and a grant from the Simons Foundation (Grant No. 230207) (E.H.L.).

<sup>1</sup> H. Araki and E. H. Lieb, "Entropy inequalities," *Commun. Math. Phys.* **18**, 160–170 (1970).

<sup>2</sup> P. Butterley, A. Sudbery, and J. Szulc, "Compatibility of subsystem states," *Found. Phys.* **36**, 83–101 (2006); e-print: [arXiv:quant-ph/0407227](https://arxiv.org/abs/quant-ph/0407227).

- <sup>3</sup>E. A. Carlen and E. H. Lieb, “Bounds for entanglement via an extension of strong subadditivity of entropy,” *Lett. Math. Phys.* **101**, 1–11 (2012); e-print: [arXiv:1203.4719v2](https://arxiv.org/abs/1203.4719v2) [quant-ph].
- <sup>4</sup>M. Christandl, B. Doron, S. Kousidis, and M. Walter, “Eigenvalue distributions of reduced density matrices,” preprint [arXiv:1204.0741](https://arxiv.org/abs/1204.0741).
- <sup>5</sup>T. Fritz and R. Chaves, “Entropic inequalities and marginal problems,” *IEEE Trans. Info. Theory* **59**(2), 803–817 (2013).
- <sup>6</sup>W. Hall, “Compatibility of subsystem states and convex geometry,” *Phys. Rev. A* **75**, 032102 (2007); e-print [arXiv:quant-ph/0610031](https://arxiv.org/abs/quant-ph/0610031).
- <sup>7</sup>See p. 33 and Eq. (4.10) in J. R. Klauder, and B.-S. Skagerstam, *Coherent States: Applications in Physics and Mathematical Physics* (World Scientific, Singapore, 1985).
- <sup>8</sup>A. Klyachko, “Quantum marginal problem and N-representability,” *J. Phys.: Conf. Ser.* **36**, 72–86 (2006).
- <sup>9</sup>S. Goldstein, J. L. Lebowitz, T. Kuna, and E. Speer, “On the extensibility of measures” (unpublished).
- <sup>10</sup>E. H. Lieb, “Convex trace functions and the Wigner-Yanase-Dyson conjecture,” *Adv. Math.* **11**, 267–288 (1973).
- <sup>11</sup>E. H. Lieb, “Some convexity and subadditivity properties of entropy,” *Bull. Am. Math. Soc.* **81**, 1–13 (1975).
- <sup>12</sup>E. H. Lieb and M. B. Ruskai, “A fundamental property of quantum mechanical entropy,” *Phys. Rev. Lett.* **30**, 434–436 (1973).
- <sup>13</sup>T. Osborne, “Entropic bounds for the quantum marginal problem,” preprint [arXiv:0806.2962](https://arxiv.org/abs/0806.2962).
- <sup>14</sup>K. R. Parthasarathy, “Extremal quantum states in coupled systems,” *Ann. I.H.P. Probab. Stat.* **41**(3), 257–268 (2005).
- <sup>15</sup>A. Perelomov, “Generalized coherent states and their applications,” *Texts and Monographs in Physics* (Springer-Verlag, Berlin, 1986).
- <sup>16</sup>O. Rudolph, “On extremal quantum states of composite systems with fixed marginals,” *J. Math. Phys.* **45**(11), 4035–4041 (2004).
- <sup>17</sup>R. Werner, “An application of Bell’s inequalities to a quantum state extension problem,” *Lett. Math. Phys.* **17**, 359–363 (1989).